


What are injection site reactions?

A common side effect that may occur with KINERET[®] treatment is a skin reaction at the injection site. These are known as **injection site reactions**.

Symptoms at the injection site may include:

- > Redness
- > Swelling
- > Bruising
- > Itching and/or stinging
- > Amyloid deposit – lump forming under the skin



Most injection site reactions are mild and usually last about **2 to 4 weeks**. However, if you are concerned about an injection site reaction or any other side effect, talk to your healthcare professional.

Please refer to the *Patient Medication Information* leaflet provided with your medication for more information.

Tips to help manage injection site reactions



- > Take KINERET out of the refrigerator and allow 30 minutes to warm to room temperature before injecting



- > Cool the site with a cold compress such as an ice pack for a few minutes – both before and after the injection



- > Rotate injection sites (i.e., change the location) to avoid soreness and the risk of a lump forming (amyloid deposit)



- > Avoid injecting into skin that is red, bruised, tattooed, tender or hard



- > Talk to your healthcare professional about over-the-counter medications, such as topical hydrocortisones or antihistamine creams, which can provide relief



Remember, there are resources available to help you with administering KINERET. Visit kineret.ca or scan to watch a demonstration on the injection procedure.

Kineret >>>>
ON TRACK[™]

If you have any questions about KINERET, contact Kineret ON TRACK[™] Program via:
☎ 1-866-204-3546, 8 am–8 pm (ET), Monday–Friday ✉ kineret@innomar-strategies.com